

Instructions Before Your FUE Procedure

TWO WEEKS PRIOR TO YOUR PROCEDURE

- **When you receive this package please call our office** to confirm your appointment and verify that we have received your laboratory test results
- If you are taking any **blood thinners**, please let the doctor know **immediately**, as they may represent a contra-indication to surgery. These medications include Plavix, Coumadin and Heparin.
- If you are taking **broad beta-blockers**, such as Inderal (Propranolol), please contact Dr. Bernstein as this medication may need to be changed to a select beta-blocker in advance of the surgery.
- Hair Length
 - **DONOR AREA:** With larger sessions of FUE (generally 1,500 or more grafts), the entire back and sides of the scalp will be shaved to 1mm in length the morning of surgery. For those having a smaller session, the doctor may shave a long strip that can be covered by the hair above it (Long-hair Robotic-FUE). **If the doctor has indicated that he may be shaving only a limited area, please leave your hair as long as possible on the back and on the sides, to cover the harvested area after your procedure.** If there is any question regarding length, it is better to leave your hair on the longer side. The doctor will trim your hair to the appropriate length the morning of surgery.
 - **RECIPIENT AREA:** On the day of surgery, it is best to have your hair the length you normally wear it. Do not shave or cut your hair in the recipient area for at least two weeks prior to your procedure. The doctor uses the hair in the recipient area (if any) to help design the hair transplant.

ONE WEEK PRIOR TO YOUR PROCEDURE

- **Do not take any B or E vitamins for one week** before your procedure this includes any multi vitamins that contain these vitamins, as these will increase bleeding. They can be resumed three days following surgery.
- **Do not take Aspirin or any other anti-inflammatory medications (NSAIDs) for one week prior to your procedure. These can be resumed three days after surgery.** NSAIDs (non-steroidal anti-inflammatory drugs include products such as Advil, Alka-Seltzer, Bufferin, Excedrin, Empirin, Nu-Prin, Motrin, Ibuprofen, etc.) If you are unsure, ask the doctor or your pharmacist.

Other prescription medications (including heart and blood pressure medication) should be taken as prescribed right up to, and including, the day of your procedure

- **Smoking:** Smoking causes constriction of blood vessels and decreased blood flow to the scalp, predominantly due to its nicotine content. The carbon monoxide in smoke

decreases the oxygen carrying capacity of the blood. These factors may contribute to poor wound healing after a hair transplant and can increase the chance of a wound infection and scarring. Smoking may also contribute to poor hair growth. The deleterious effects of smoking wear off slowly when one abstains, particularly in chronic smokers, so that smoking puts one at risk to poor healing even after smoking is stopped for weeks or even months. Although it is not known exactly how long one should avoid smoking before and after a hair transplant a common recommendation is to abstain from **one week prior** to surgery until **two weeks** after the procedure.

THREE DAYS PRIOR TO YOUR PROCEDURE

- **Do not drink any alcoholic beverages** for at least **three days** prior to your procedure.

ONE DAY PRIOR TO YOUR PROCEDURE

- It is difficult for the ARTAS robot to identify white or light-blond hair. **If you have white or blond hair, please dye your hair light brown (or darker) the day before your procedure, as this will make it easier for the robot to visualize your hair.** This should be repeated the morning of your procedure (when possible). If the product comes with a stain remover, try to remove the dye-stain from your scalp.

THE DAY OF YOUR PROCEDURE

- **Shower the morning of your procedure and wash your hair very thoroughly using your regular shampoo.** Do not apply sprays, gels, or any other styling products to your hair. If you wear a hair system, please remove it prior to shampooing and do not wear it before your procedure.
- If time permits, the morning of surgery repeat hair dying as per above.
- **Wear loose, comfortable clothing which does not need to be pulled on over your head.** This will prevent dislodging your bandage or damaging your grafts immediately after surgery.
- Please do not wear any cologne or perfume the day of your procedure.
- **Eat breakfast the morning of your procedure.** We will provide lunch. If your surgery is scheduled for the afternoon, have a light lunch before you arrive.
- **Do not drink coffee or any caffeinated beverages on the day of your procedure.** These substances may increase bleeding and your sensitivity to medications used during the procedure.
- You may be receiving medications during the procedure that can make you drowsy. Because of this, **you cannot drive yourself home following your procedure.** Please arrange to have a responsible person drive you to your destination. If you are unable to have someone drive you home after your procedure and you live near our office, we will call a taxi for you and you will be responsible for the charges.
- Please do not bring any valuables to the office on the day of surgery. This includes laptop computers.

- You will be going home with a headband-type bandage covered by a bandana, so it will not be visible. The bandages should be removed just before showering the next morning. After the first shower, no more bandages are need.

If you have any questions, please contact us at 212-826-2400.